

Horseshoe Crab Spawning Activity in Delaware Bay: 1999 – 2013

Report to the Atlantic States Marine Fisheries Commission's Horseshoe Crab Technical Committee

Jordan Zimmerman, Delaware Division of Fish and Wildlife,
89 Kings Highway, Dover, DE 19901
(302) 735-2976 / jordan.zimmerman@state.de.us

Edward Hale, Delaware Division of Fish and Wildlife,
89 Kings Highway, Dover, DE 19901
(302) 739-2969 / ed.hale@state.de.us

David Smith, USGS – Leetown Science Center,
11649 Leetown Road, Kearneysville, WV 25443
(304) 724-4467 / drsmith@usgs.gov

Sherry Bennett, New Jersey Division of Fish, Game and Wildlife,
PO Box 418, Port Republic, NJ 08241

July 16, 2014

Summary

- This annual report marks the fifteenth year that the Delaware Bay Horseshoe Crab Spawning Survey has been implemented in a standardized manner throughout May and June in the Delaware Bay.
- Annual coefficients of variation for estimates of female spawning activity were below 14% for the entire series and remained at or below 10% for the last ten years. Annual coefficients of variation for estimates of male spawning activity were below 20% for the entire series.
- Female spawning activity in 2013 peaked during the second lunar period sampled (May 23 – May 27).
- The proportion of female spawning activity observed in May 2013 in New Jersey (62%) was below the 15 year average for that state (66%) while Delaware (71%) was well above average (59%).
- Percent of female spawning that occurred in May was associated with water temperature (correlations were 0.53 and 0.62 for DE and NJ, respectively).
- Bay-wide female spawning activity over the past 15 years showed no significant trend; though, the slope was slightly negative (Slope = -0.01, SE = 0.01, 90% CI = -0.02 to 0.01, P = 0.29).
- No significant trends in state-specific female spawning were detected; though, the slope in both Delaware and New Jersey was negative. (DE: slope = -0.01, SE = 0.01, 90% CI = -0.02 to 0.01, P = 0.29; NJ: slope = -0.01, SE = 0.01, 90% CI = -0.02 to 0.01, P = 0.84).
- Bay-wide male spawning activity showed no significant trend from 1999 through 2013; though, the slope was positive (Slope = 0.05, SE = 0.05, 90% CI = -0.03 to 0.12, P = 0.30).

- No significant trends in state-specific male spawning were detected; although, the slopes in both states were positive (DE: slope = 0.02, SE = 0.05, 90% CI = -0.02 to 0.01, P = 0.72; NJ; slope = 0.06, SE = 0.06, 90% CI = -0.02 to 0.01, P = 0.30).
- Sex ratios (M:F) during the 15 year time series ranged from 3.1 to 5.2.

Introduction

The Atlantic States Marine Fisheries Commission's (ASMFC) Interstate Fishery Management Plan for Horseshoe Crab (ASMFC 1998) required that the states of Delaware, Maryland and New Jersey implement pilot horseshoe crab spawning surveys based on "standardized and statistically robust methodologies". In January 1999, the ASMFC convened a workshop that established a framework for such surveys in the Mid-Atlantic region. The framework built upon existing horseshoe crab spawning survey efforts by Finn et al. (1991) and Maio (1998). Using funds from the U.S. Geological Survey's (USGS) State Partnership Program, a comprehensive pilot study was designed and implemented in Delaware Bay during the spring of 1999 (Smith et al. 2002). The U.S. Fish and Wildlife Service provided further funding in 2000 to continue the survey in its present form, and the Delaware Division of Fish and Wildlife (DE DFW) provided funding in subsequent years using Atlantic Coastal Fisheries Cooperative Management Act funds. The survey has been shown to provide levels of spatial and temporal coverage essential for understanding trends in spawning activity (Smith and Michels 2006).

The survey is an excellent example of state, federal, non-governmental organization (NGO), corporate and citizen cooperation. Survey coordination is contracted through Limuli Labs. Data entry is completed by staff from the New Jersey Department of Environmental Protection; USGS and DE DFW staff oversee data analysis and report preparation. The vast sampling effort is conducted by a large contingent of dedicated private citizens, state and federal agencies, corporations, and NGO's.

This report is a continuation of a series of statistical reports on the survey and is meant to compliment the ongoing series of reports issued by the survey coordinators, Ms. Benjie Swan and Dr. William Hall in cooperation with Dr. Carl N. Shuster Jr.

Survey Objectives

The Delaware Bay Horseshoe Crab Spawning Survey has several important objectives:

- 1) Provide a reliable index of spawning activity to monitor the temporal and spatial distribution of horseshoe crab spawning activity for comparing bay-wide spawning among years, beach-level spawning within Delaware Bay, and distributions of spawning horseshoe crabs and shorebirds;
- 2) Increase our understanding of the relationship between environmental factors (tidal height, wave height, and water temperature) and spawning activity; and
- 3) Promote public awareness of the central role of horseshoe crabs in shorebird population dynamics, Atlantic coast fisheries, and human health through the production of *Limulus* ameobocyte lysate (LAL).

Data Availability

The spawning survey database was converted to MS ACCESS in 2004. A visual basic program was also developed by USGS to calculate estimates of spawning activity in tabular and graphic form. The conversion process revealed a number of errors that were corrected and detailed in Smith and Bennett (2005). The overall patterns of spawning activity were largely unaffected by these corrections. Beginning in 2010, the previous software was no longer compatible with updated Windows OS, so the SPAWNr program was developed by Dr. David Smith (USGS) to calculate estimates. Data used in this report (both estimates and raw data) and the software used to calculate estimates are available by request. In 2014, it was discovered that the calculation of the reported standard deviation for bay-wide mean spawning activity was denoted incorrectly as the standard error of such data. This has been corrected and standard deviation will continue to be reported in the future.

Summary Results

Sampling in 2013 was conducted during twelve nighttime high tides from 7 May through 25 June. Twenty-five beaches were sampled in the Delaware Estuary – 13 in Delaware and 12 in New Jersey. The total number of tides sampled over the season was 275, with 25 sampling events missed or canceled (Table 1). Twelve of the 25 missed sampling events occurred during the second and third lunar periods when spawning horseshoe crabs were most abundant.

Table 1. Beaches sampled in the 2013 Delaware Bay Horseshoe Crab Spawning Survey.

Beach	May						June					
	7	9	11	23	25	27	6	8	10	21	23	25
<u>Delaware</u>												
Woodland										Partial Count		
Pickering												
Kitts Hummock												
Ted Harvey				Partial Count								
N. Bowers												
S. Bowers												
Bennetts Pier				Partial Count								
Big Stone												
Slaughter		Partial Count										
Fowler												
Prime Hook				Partial Count								
Broadkill												
Cape Henlopen										Partial Count		
<u>New Jersey</u>												
Fortescue												
Reeds												
Gandy's		Partial Count										
Kimbles												
Higbees												
Pierces Point												
Higs												
Norburys												
S. Cape Shore Lab				Partial Count								
Villas												
N. Cape May				Partial Count								
Townbank												

Sampled

- Sampled
- Partial Count

Not Sampled

- No Access / Flooding
- Weather
- No Surveyors
- No data / Other

Temporal Spawning Distribution

Horseshoe crab phenology is an important factor to examine as it gives an indication of the timing of potential food availability to migratory shorebirds. The time of spawning may also affect the survival of egg, larvae and juvenile stages.

State-specific female spawning activity peaked in New Jersey and Delaware in the first (May 7 - 11) and second (May 23 - 27) lunar periods, respectively (Figure 1). Sixty-two percent (62%) of the annual female spawning activity in New Jersey and 71% of the annual female spawning activity in Delaware was observed in May (Table 2). The proportion of annual state-specific spawning activity that occurred in May was higher in New Jersey than Delaware for all but three years of the 15 year survey.

Bay-wide female spawning activity peaked in the second lunar period in 2013 (Table 3). This is the tenth year of the 15 year survey that the second lunar period in May has accounted for the highest spawning activity of the survey. This period is critical to shorebird foraging as it coincides with peak stopover period for migratory shorebirds in Delaware Bay.




Figure 1. Temporal distribution of female horseshoe crab spawning activity in the Delaware Bay by state for the years 2009-2013. Lunar periods are defined as a 5 day period (sampled day of lunar event and 2 days before and 2 days after) around the new or full moons in May and June.

Table 2. Summary statistics reflecting the timing of female horseshoe crab spawning in Delaware and New Jersey and average May water temperatures. Water temperatures were recorded at the National Ocean Service station at Lewes, DE.

	Delaware		New Jersey		Average daily water temp. in May (C)
	Dates of Peak Female Spawning	% of Female Spawning in May	Dates of Peak Female Spawning	% of Female Spawning in May	
1999	28 May - 1 June	77	28 May - 1 June	93	16.2
2000	16 May - 18 May	54	16 May - 18 May	64	15.6
2001	3 June - 7 June	47	5 May - 9 May	76	16.0
2002	24 May - 28 May	73	24 May - 28 May	78	16.7
2003	29 May - 2 June	47	29 May - 2 June	56	13.4
2004	17 May - 21 May	76	17 May - 21 May	85	15.7
2005	4 June - 8 June	18	4 June - 8 June	30	13.7
2006	25 May - 29 May	77	25 May - 29 May	85	16.3
2007	30 May - 3 June	42	30 May - 3 June	45	15.4
2008	1 June - 5 June	43	1 June - 5 June	26	15.2
2009	22 May - 26 May	59	22 May - 26 May	66	15.5
2010	12 May - 16 May	82	25 May - 29 May	88	15.6
2011	30 May - 3 June	52	30 May - 3 June	44	16.0
2012	2 June - 6 June	64	18 May - 22 May	92	17.8
2013	23 May - 27 May	71	7 May - 11 May	62	15.3

Table 3. Baywide horseshoe crab spawning activity, expressed as mean number of spawning female crabs per m², by lunar period for the years 1999 to 2013.

Year	0	1	2	3	4	5
1999		0.86	1.58	0.32	0.15	
2000		0.92	1.23	0.91	0.62	
2001		0.77	0.96	0.76	0.42	
2002		0.92	1.81	0.71	0.14	
2003		0.04	0.17	1.51	1.13	0.46
2004		0.56	1.91	0.30	0.30	
2005		0.12	0.67	2.00	0.36	
2006		1.39	1.85	0.61	0.11	
2007		0.17	1.34	1.61	0.38	
2008		0.78	0.17	1.49	0.22	
2009		0.67	1.84	0.96	0.48	
2010		1.26	1.52	0.42	0.06	
2011		0.46	0.92	1.00	0.21	
2012	0.18*	0.71	1.02	0.43	0.14	
2013		0.86	1.30	0.68	0.53	

*denotes partial survey

Water temperature likely influences the time of spawning (Smith and Michels 2006). There was a strong association between average May water temperatures recorded at Lewes, DE and the percentage of state-specific female spawning activity in May ($r_{DE} = 0.53$, $P_{DE} = 0.04$; $r_{NJ} = 0.62$, $P_{NJ} = 0.01$; Figure 2). Delayed spawning in 2003 and 2005 was likely related to water temperatures, as temperatures were not consistently above 15 °C until late May or early June at Lewes, DE (Appendix I). Reduced spawning activity noted during the second lunar sampling period in 2008 coincided with a severe nor'easter that depressed water temperatures. In 2013, water temperatures did not remain consistently above 15°C until May 17, well after the first lunar period was sampled in the spawning survey.




Figure 2. Percent of female horseshoe crab spawning occurring in May by state versus May average daily water temperatures. Water temperatures were recorded by the National Ocean Service at Lewes, DE Station ID 8557380.

State-specific Spawning Activity

Although index values differ by state (Table 4; Figure 3), the trend from the index of female spawning activity in both states exhibited a slightly negative slope, though not significant (DE Slope = -0.007, SE = 0.007, P = 0.30; NJ Slope = -0.003, SE = 0.01, P = 0.85).

Trends in male spawning activity were similar between states (Table 5; Figure 4). The index of male spawning activity in both states exhibited a positive slope, though not significant (DE Slope = 0.02, SE = 0.05, P = 0.72; NJ Slope = 0.07, SE = 0.06, P = 0.30).

Table 4. Indices of female horseshoe crab spawning activity (IFSA), expressed as the mean number of female crabs per m^2 , by state from 1999 to 2013.

Year	Delaware			New Jersey		
	IFSA	90% CI	Beaches Surveyed	IFSA	90% CI	Beaches Surveyed
1999	0.93	0.67, 1.29	8	0.61	0.47, 0.80	9
2000	1.02	0.72, 1.45	11	0.80	0.67, 0.96	11
2001	0.82	0.63, 1.08	12	0.64	0.51, 0.80	10
2002	0.76	0.61, 0.94	13	1.09	0.92, 1.30	10
2003	0.81	0.64, 1.03	13	0.83	0.76, 0.91	10
2004	0.76	0.62, 0.93	13	0.78	0.68, 0.89	12
2005	0.65	0.53, 0.80	13	0.99	0.84, 1.16	12
2006	0.81	0.67, 0.98	13	1.17	1.03, 1.33	11
2007	0.96	0.79, 1.15	13	0.82	0.68, 0.99	11
2008	0.78	0.63, 0.96	13	0.57	0.49, 0.67	12
2009	0.73	0.60, 0.90	13	1.26	1.11, 1.42	13
2010	0.79	0.64, 0.99	13	0.81	0.68, 0.96	12
2011	0.71	0.59, 0.85	13	0.56	0.48, 0.65	12
2012	0.45	0.33, 0.62	13	0.68	0.55, 0.83	12
2013	0.93	0.72, 1.20	13	0.73	0.61, 0.87	12

Delaware Female




Figure 3a. Indices of female horseshoe crab spawning activity (IFSA), expressed as the mean number of female crabs per m^2 , for the state of Delaware. Error bars are 90% confidence intervals.




Figure 3b. Indices of female horseshoe crab spawning activity (IFSA), expressed as the mean number of female crabs per m^2 , for the state of New Jersey. Error bars are 90% confidence intervals.

DRAFT

Table 5. Indices of male horseshoe crab spawning activity (IMSA), expressed as the mean number of male crabs per m² per night, by state from 1999 to 2013.

Year	Delaware			New Jersey		
	IMSA	90% CI	Beaches Surveyed	IMSA	90% CI	Beaches Surveyed
1999	3.78	2.65, 5.37	8	1.82	1.24, 2.65	9
2000	3.93	2.76, 5.60	11	2.00	1.55, 2.59	11
2001	2.76	2.02, 3.76	12	2.01	1.50, 2.69	10
2002	2.74	2.13, 3.52	13	3.43	2.91, 4.06	10
2003	2.90	2.23, 3.77	13	2.98	2.67, 3.33	10
2004	2.85	2.27, 3.59	13	3.07	2.64, 3.57	12
2005	2.49	1.99, 3.11	13	4.00	3.30, 4.85	12
2006	3.80	3.03, 4.75	13	4.45	3.84, 5.15	11
2007	4.64	3.81, 5.66	13	4.00	3.22, 4.97	11
2008	4.03	3.16, 5.14	13	2.23	1.86, 2.69	12
2009	3.87	3.08, 4.87	13	5.46	4.74, 6.30	13
2010	3.48	2.77, 4.38	13	3.31	2.75, 3.99	12
2011	4.36	3.49, 5.45	13	2.24	1.93, 2.61	12
2012	2.10	1.48, 3.01	13	2.77	2.15, 3.57	12
2013	3.38	2.61, 4.39	13	2.88	2.35, 3.54	12

Delaware Male




Figure 4a. Indices of male horseshoe crab spawning activity (IMSA), expressed as the mean number of male crabs per m², for the state of Delaware. Error bars are 90% confidence intervals.




Figure 4b. Indices of male horseshoe crab spawning activity (IMSA), expressed as the mean number of male crabs per m^2 , for the state of New Jersey. Error bars are 90% confidence intervals.

Bay-wide Spawning Activity - Females

Trends in state-specific female spawning activity were compensatory, as no change in bay-wide spawning activity was detected (Figure 5; Table 6). The regression slope was close to zero (Slope = -0.01, SE = 0.01, 90% CI = -0.02 to 0.005, P = 0.29). Coefficients of variation were below 14% over the entire survey period and at or below 10% since 2002. Female spawning activity by beach for all years is provided in Appendix II. Smith and Robinson (2014) recently used mixed-model trend regression to evaluate beach level trends in spawning density. Their results indicate that while concentrations at primary spawning beaches tend to be stabilizing, higher numbers of spawning females have become more numerous among ancillary Delaware Bay beaches.

Baywide Female




Figure 5. Index of female horseshoe crab spawning activity (IFSA) for the Delaware Bay from 1999 to 2013. Error bars are 90% confidence intervals.

Table 6. Indices of bay-wide male and female horseshoe crab spawning activity (ISA), number of beaches surveyed, standard deviations (SD), coefficient of variations (CV), 90% confidence intervals (CI) and sex ratio for the Delaware Bay from 1999 to 2013.

Year	Beaches Surveyed	Male				Female				Annual Sex Ratio (M:F)
		ISA	90% CI	SD	CV (%)	ISA	90% CI	SD	CV (%)	
1999	17	2.50	1.86, 3.37	0.45	18	0.77	0.62, 0.97	0.10	13	3.2
2000	22	2.96	2.31, 3.80	0.45	15	0.91	0.74, 1.13	0.12	13	3.2
2001	22	2.37	1.91, 2.95	0.31	13	0.75	0.63, 0.90	0.08	10	3.1
2002	23	2.86	2.45, 3.34	0.27	9	0.91	0.79, 1.04	0.07	8	3.1
2003	23	2.89	2.50, 3.33	0.25	9	0.80	0.71, 0.91	0.06	8	3.6
2004	24	2.93	2.55, 3.36	0.24	8	0.77	0.68, 0.87	0.06	7	3.8
2005	23	3.23	2.79, 3.74	0.29	9	0.82	0.72, 0.93	0.07	9	3.9
2006	24	3.99	3.49, 4.56	0.33	8	0.99	0.89, 1.10	0.07	7	4.0
2007	24	4.22	3.63, 4.90	0.38	9	0.89	0.78, 1.01	0.07	8	4.7
2008	25	2.30	1.83, 2.90	0.32	14	0.68	0.59, 0.78	0.06	9	3.4
2009	26	4.67	4.11, 5.29	0.36	8	1.00	0.89, 1.11	0.06	6	4.7
2010	25	3.39	2.93, 3.94	0.31	9	0.80	0.70, 0.92	0.07	8	4.2
2011	25	3.31	2.83, 3.87	0.31	10	0.64	0.57, 0.72	0.05	7	5.2
2012	25	2.44	1.97, 3.01	0.31	13	0.56	0.47, 0.67	0.06	10	4.4
2013	25	3.14	2.65, 3.71	0.32	10	0.83	0.71, 0.98	0.08	10	3.8

Survey Sex Ratios

Current horseshoe crab harvest management strategies in the Delaware Bay area favor the harvest of male crabs. Concern was expressed that these strategies may cause spawning sex ratios (M:F) to drop and negatively affect spawning and egg fertilization. Annual sex ratio has ranged from 3.1 to 5.2 over the course of the survey. M:F ratio in 2013 (3.8) was below four for the first time since 2008 (Table 6). Bay-wide Spawning Activity - Males

Sex-specific harvest requirements contained in Addendum IV to the Interstate Fishery Management Plan for Horseshoe Crab (ASMFC 2006) for Delaware and New Jersey (specifically a male-only harvest) prompted an examination of bay-wide male spawning abundance. Male spawning activity increased slightly, though not significantly (Slope = 0.05, SE = 0.04, 90% CI = -0.02 to 0.12, P = 0.30) from 1999 to 2013 (Figure 6; Table 6). Coefficients of variation for the male component of the survey were below 20% for the entire sampling period.





Figure 6. Index of male horseshoe crab spawning activity (IMSA) for the Delaware Bay from 1999 to 2013. Error bars are 90% confidence intervals.

Literature Cited

- Atlantic States Marine Fisheries Commission (ASMFC). 1998. Interstate fishery management plan for horseshoe crab. Fishery Management Report No. 32, Atlantic States Marine Fisheries Commission, Washington D.C.
- Atlantic States Marine Fisheries Commission (ASMFC). 2006. Addendum IV to the Interstate Fishery Management Plan for Horseshoe Crab. Fishery Management Report No. 32d, Atlantic States Marine Fisheries Commission, Washington D.C.
- Carmichael, R.H., D. Rutecki and I. Valiela. 2003. Abundance and population structure of the Atlantic horseshoe crab *Limulus polyphemus* in Pleasant Bay, Cape Cod. *Marine Ecol. Prog. Ser.* 246:225-239.
- Finn, J.J., C.N. Shuster, Jr. and B.L. Swan. 1991. *Limulus* spawning activity on Delaware Bay shores 1990. Finn-Tech Inc., Cape May Courthouse, NJ.
- Maio, K.J. 1998. Quantitative description of the temporal and spatial patterns in spawning activity of the horseshoe crab, *Limulus polyphemus*. Master's thesis. University of Maryland Eastern Shore, Princess Anne, Maryland.
- Smith, D.R., P.S. Pooler, B.L. Swan, S.F. Michels, W.R. Hall, P.J. Himchak, and M.J. Millard. 2002. Spatial and temporal distribution of horseshoe crab (*Limulus polyphemus*) spawning in Delaware Bay: implications for monitoring. *Estuaries* 25(1):115-125.
- Smith D.R. and S. Bennett. 2005. Horseshoe crab spawning activity in Delaware Bay: 1999 – 2004. Report to the ASMFC Horseshoe Crab Management Board. Unpublished.
- Smith, D.R., and S.F. Michels. 2006. Seeing the elephant: importance of spatial and temporal coverage in a large-scale volunteer-based program to monitor horseshoe crabs. *Fisheries* 31(10):485-491.
- Smith, D.R., and T.J. Robinson. 2014. Delaware Bay horseshoe crab spawning activity after harvest reduction based on mixed-model analyses. In review.
- Zimmerman, J., S. Michels, D. Smith, and S. Bennett. 2012. Horseshoe crab spawning activity In Delaware Bay: 1999 – 2012. Unpublished report to the ASMFC Horseshoe Crab Technical Committee.

APPENDIX I. Water temperature data from Lewes, DE (Station Identification Number 8557380; Latitude 38° 46.9' N / Longitude 75° 7.2' W) for the 2013. Source: Center for Operational Oceanographic Products and Services (CO-OPS).



APPENDIX II. Index of female spawning horseshoe crabs abundance, expressed as the mean number of female crabs per m² per night, for Delaware Bay beaches surveyed from 1999 to 2013.

State	Beach	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
DE	Bennetts Pier		0.22	0.64	0.47	0.28	0.55	0.70	0.61	0.55	0.37	0.64	0.22	0.28	0.32	0.13
DE	Big Stone	0.75	0.73	0.86	0.63	0.64	0.76	0.81	1.09	1.35	0.71	0.79	0.67	0.86	0.54	0.54
DE	Broadkill	0.32	0.06	0.12	0.13	0.21	0.17	0.19	0.12	0.18	0.55	0.23	0.49	0.57	0.22	0.57
DE	Cape Henlopen				0.09	0.18	0.13	0.27	0.10	0.06	0.30	0.26	0.32	0.42	0.30	0.42
DE	Fowlers	0.78	0.49	0.70	0.24	0.45	0.61	0.21	0.41	0.50	0.53	0.21	0.42	0.13	0.06	0.31
DE	Kitts Hummock	2.15	2.58	2.35	1.47	1.55	1.24	1.42	1.72	1.44	1.23	1.48	1.30	1.27	0.85	1.91
DE	Lewes				0.08											
DE	North Bowers	0.88	1.18	1.04	1.21	0.98	0.50	0.60	0.75	1.11	0.36	0.69	0.75	0.49	0.43	1.08
DE	Pickering		3.30	1.62	1.70	1.64	1.64	1.47	1.49	1.64	1.99	1.67	1.87	1.14	1.42	2.55
DE	Prime Hook	0.60	0.19	0.44	0.59	0.47	0.76	0.65	0.73	1.11	0.92	0.61	0.92	1.03	0.26	1.12
DE	Slaughter	1.62	1.33	1.10	0.73	1.65	1.52	0.68	1.04	1.24	1.10	0.72	0.75	1.14	0.47	1.47
DE	South Bowers		0.92	0.84	1.13	0.47	0.48	0.63	0.72	1.30	0.57	1.02	0.50	0.58	0.54	0.66
DE	Ted Harvey				1.44	1.99	1.52	0.82	1.46	1.93	1.47	1.19	1.34	1.35	1.23	2.13
DE	Woodland	0.14	0.10	0.03	0.08	0.01	0.00	0.01	0.27	0.03	0.00	0.02	0.16	0.01	0.08	0.01
NJ	East Point		0.35													
NJ	Fortescue	0.25				0.42	0.54	0.58	0.65	0.16	0.33	0.44	0.34	0.34	0.73	0.93
NJ	Gandys	0.40	0.39	0.45	1.41	0.55	0.82	0.88	1.17	0.83	0.30	1.31	1.24	0.25	1.50	1.08
NJ	Higbees		0.04					0.14			0.03	0.14		0.42	0.06	0.07
NJ	Highs Beach	0.79	0.96	0.80	0.47	0.53	0.70	0.76	0.69	0.75	0.46	0.73	0.56	0.61	0.68	0.71
NJ	Kimbles	0.71	0.85	0.48	0.50	0.50	0.41					0.82	0.51	0.33	0.93	0.49
NJ	Norburys			0.46	0.62	0.54	0.67	0.94	0.69	0.43	0.41	1.14	0.68	0.71	0.78	0.76
NJ	North Cape May	0.23	0.05	0.09	0.08	0.12	0.02	0.12	0.02	0.04	0.03	0.08	0.02	0.24	0.03	0.06
NJ	Pierces Point		0.61		0.67	0.73	0.96	0.83	0.74	0.94	0.71	1.27	1.11	0.96	1.13	1.19
NJ	Raybins	0.03														
NJ	Reeds	0.38	0.65	0.40	0.88	0.82	0.42	0.24	0.97	0.31	0.34	1.07	0.57	0.82	0.86	0.95
NJ	Sea Breeze	0.09	0.11	0.30	1.63	0.39	0.43	0.21	0.85	0.93	0.67	0.77	1.02			
NJ	Cape Shore Lab	1.25	1.33	1.28	0.69	0.63	0.90	1.17	0.82	1.26	0.39	1.11	0.79	0.80	1.19	0.80
NJ	Sunset			0.11					0.01	0.00	0.01	0.16				
NJ	Townbank			0.74	0.40	0.46	0.20			0.29			0.31	0.39	0.24	0.52
NJ	Villas							0.71	0.48		0.34	0.64	0.41	0.53	0.24	0.35